

THE MELLIN TRANSFORM OF THE SQUARE OF RIEMANN'S ZETA-FUNCTION

ALEKSANDAR IVIĆ

Abstract. Let $\mathcal{Z}_1(s) = \int_1^\infty |\zeta(\frac{1}{2} + ix)|^2 x^{-s} dx$ ($\sigma = \Re s > 1$). A result concerning analytic continuation of $\mathcal{Z}_1(s)$ to $\mathbb{C}$ is proved, and also a result relating the order of $\mathcal{Z}_1(\sigma + it)$ ($\frac{1}{2} \leq \sigma \leq 1, t \geq t_0$) to the order of $\mathcal{Z}_1(\frac{1}{2} + it)$.

1. Introduction

Let $\mathcal{Z}_k(s)$, the (modified) Mellin transform of $|\zeta(\frac{1}{2} + ix)|^{2k}$, denote the analytic continuation of the function defined initially by

$$(1.1) \quad \mathcal{Z}_k(s) := \int_1^\infty |\zeta(\frac{1}{2} + ix)|^{2k} x^{-s} dx \quad (k \in \mathbb{N}, \sigma = \Re s > c(k) (> 1)).$$

This function, when $k = 2$, was introduced by Y. Motohashi [15] (see also [16]), and its properties were further studied in [10] and [11]. The latter work also contains some results on the function $\mathcal{Z}_1(s)$, which is the principal object of the study in this paper. It was shown that $\mathcal{Z}_1(s)$ is regular for $\sigma > -3/4$, except for a double pole at $s = 1$. The principal part of the Laurent expansion of $\mathcal{Z}_1(s)$ at $s = 1$ is

$$(1.2) \quad \frac{1}{(s-1)^2} + \frac{2\gamma - \log(2\pi)}{s-1},$$

where $\gamma = -\Gamma'(1) = 0.577215 \dots$ is Euler's constant. M. Jutila [13] continued the study of $\mathcal{Z}_1(s)$ and proved that $\mathcal{Z}_1(s)$ continues meromorphically to $\mathbb{C}$, having only a double pole at $s = 1$ and at most double poles for $s = -1, -2, \dots$.

The object of this note is to prove some new results on $\mathcal{Z}_1(s)$. First we make more precise Jutila's result on the analytic continuation of $\mathcal{Z}_1(s)$. We have the following

THEOREM 1. *The function $\mathcal{Z}_1(s)$ continues meromorphically to $\mathbb{C}$, having only a double pole at $s = 1$, and at most simple poles at $s = -1, -3, \dots$. The principal part of its Laurent expansion at $s = 1$ is given by (1.2).*

Our second aim is to prove an order result for $\mathcal{Z}_1(s)$. This is a result M. Jutila mentioned in [13], but the term that I initially claimed, namely $t^{1-2\sigma+\varepsilon}$ is too optimistic. It appears that what can be proved is contained in

THEOREM 2. *We have, for $\frac{1}{2} \leq \sigma \leq 1, t \geq t_0 > 0$,*

$$(1.3) \quad \mathcal{Z}_1(\sigma + it) \ll_\varepsilon t^{\frac{1}{2}-\sigma+\varepsilon} \max_{t-t^\varepsilon \leq v \leq t+t^\varepsilon} |\mathcal{Z}_1(\frac{1}{2} + iv)| + (t^{\frac{9-16\sigma}{7}} + t^{-1}) \log t.$$

Corollary (see M. Jutila [13]). For $\frac{1}{2} \leq \sigma \leq 1$, $t \geq t_0$ we have

$$(1.4) \quad \mathcal{Z}_1(\sigma + it) \ll_{\varepsilon} t^{\frac{5}{6} - \sigma + \varepsilon}.$$

The bound $\mathcal{Z}_1(\frac{1}{2} + it) \ll_{\varepsilon} t^{1/3 + \varepsilon}$ was mentioned in [11], and its proof was elaborated by M. Jutila in [13]. If one inserts this bound in (1.3), then (1.4) follows immediately. We note that here and later ε denotes arbitrarily small constants, not necessarily the same ones at each occurrence. It can be also proved that, for any given $\varepsilon > 0$,

$$(1.5) \quad \int_1^T |\mathcal{Z}_k(\sigma + it)|^2 dt \gg_{\varepsilon} T^{2-2\sigma-\varepsilon} \quad (k = 1, 2; \frac{1}{2} < \sigma < 1).$$

The bound (1.5) for $k = 2$ appeared in my paper [10], and the proof of the bound when $k = 1$ is on similar lines, so that the details will not be given here. It is plausible that

$$(1.6) \quad \max_{t-t^{\varepsilon} \leq v \leq t+t^{\varepsilon}} |\mathcal{Z}_1(\frac{1}{2} + iv)| \gg 1$$

holds, but at present I am unable to prove (1.6).

2. Proof of Theorem 1

Let

$$(2.1) \quad L_k(s) := \int_0^{\infty} |\zeta(\frac{1}{2} + ix)|^{2k} e^{-sx} dx \quad (\sigma = \Re s > 0, k \in \mathbb{N})$$

be the Laplace transform of $|\zeta(\frac{1}{2} + ix)|^{2k}$. We are interested in using the expression for $L_1(s)$ when $s = 1/T$, $T \rightarrow \infty$. Such a formula has been known for a long time and is due to H. Kober [14] (see also a proof in [17, Chapter 9]). The functions $L_k(s)$ when $k = 1, 2$ were studied by F.V. Atkinson [1], [2]. The function $L_2(s)$ was considered by the author [8], [9], and the approach to mean value of Dirichlet series by Laplace transforms by M. Jutila [12]. Kober's result on $L_1(s)$ is that, for any integer $N \geq 0$,

$$(2.2) \quad L_1(\sigma) = \frac{\gamma - \log(2\pi\sigma)}{2 \sin \frac{\sigma}{2}} + \sum_{n=0}^N d_n \sigma^n + O_N(\sigma^{N+1}) \quad (\sigma \rightarrow 0+),$$

where d_n are suitable constants. Note that for $\sigma = 1/T$, $T \rightarrow \infty$ we have

$$\begin{aligned} \frac{\gamma - \log(\frac{2\pi}{T})}{2 \sin(\frac{1}{2T})} &= \frac{\log(\frac{T}{2\pi}) + \gamma}{\frac{1}{T} - \frac{1}{24T^3} + \frac{1}{16.5!T^5} - \dots} \\ &= \left(\log\left(\frac{T}{2\pi}\right) + \gamma \right) \sum_{n=0}^{\infty} c_n T^{1-2n} \end{aligned}$$

with the coefficients c_n that may be explicitly evaluated. Therefore for any integer $N \geq 0$

$$(2.3) \quad L_1\left(\frac{1}{T}\right) = \left(\log\left(\frac{T}{2\pi}\right) + \gamma\right) \sum_{n=0}^N a_n T^{1-2n} + \sum_{n=0}^N b_n T^{-2n} + O_N(T^{-1-2N} \log T)$$

with the coefficients a_n, b_n that may be explicitly evaluated. In particular, we have

$$(2.4) \quad a_0 = 1, \quad b_0 = \pi.$$

It is clear from Kober's formula (2.2) that $a_0 = 1$. To see that $b_0 = \pi$ one can use the work of Hafner–Ivić [4] (this was mentioned by Conrey et al. [3]), which will be stated now in detail, as it will be needed also for the proof of Theorem 2. Let, as usual, for $T \geq 0$,

$$E(T) = \int_0^T |\zeta(\tfrac{1}{2} + it)|^2 dt - T \left(\log\left(\frac{T}{2\pi}\right) + 2\gamma - 1 \right) \quad (E(0) = 0)$$

denote the error term in the mean square formula for $|\zeta(\tfrac{1}{2} + it)|$. Let further

$$(2.5) \quad G(T) := \int_0^T E(t) dt - \pi T, \quad G_1(T) := \int_0^T G(t) dt.$$

Then Hafner–Ivić [4] proved (see also [6])

$$(2.6) \quad G(T) = S_1(T; N) - S_2(T; N) + O(T^{1/4})$$

with

$$S_1(T; N) = 2^{-3/2} \sum_{n \leq N} (-1)^n d(n) n^{-1/2} \left(\operatorname{arsinh} \sqrt{\frac{\pi n}{2T}} \right)^{-2} \left(\frac{T}{2\pi n} + \frac{1}{4} \right)^{-1/4} \sin(f(T, n)),$$

$$S_2(T; N) = \sum_{n \leq N'} d(n) n^{-1/2} \left(\log \frac{T}{2\pi n} \right)^{-2} \sin(g(T, n)),$$

$$f(T, n) = 2T \operatorname{arsinh} \sqrt{\frac{\pi n}{2T}} + \sqrt{2\pi n T + \pi^2 n^2} - \frac{\pi}{4},$$

$$g(T, n) = T \log \left(\frac{T}{2\pi n} \right) - T + \frac{\pi}{4}, \quad \operatorname{arsinh} x = \log(x + \sqrt{x^2 + 1}),$$

$$AT < N < A'T \quad (0 < A < A' \text{ constants}), \quad N' = \frac{T}{2\pi} + \frac{N}{2} - \sqrt{\frac{N^2}{4} + \frac{NT}{2\pi}}.$$

We use Taylor's formula (see [7, Lemma 3.2]) to simplify (2.6). Then we obtain

$$G(T) = 2^{-1/4} \pi^{-3/4} T^{3/4} \sum_{n=1}^{\infty} (-1)^n d(n) n^{-5/4} \sin(\sqrt{8\pi n T} - \frac{\pi}{4}) + O(T^{2/3} \log T),$$

so that

$$(2.7) \quad G(T) = O(T^{3/4}), \quad G(T) = \Omega_{\pm}(T^{3/4}).$$

We use (2.6) with $T = t, N = T, T \leq t \leq 2T$ and apply the first derivative test ([5, Lemma 2.1]) to deduce that

$$\int_T^{2T} G(t) dt \ll T^{5/4},$$

since the term $O(1)$ in [5, eq. (15.29)] in Atkinson's formula is in fact $O(1/T)$. Hence

$$(2.8) \quad G_1(T) = \int_0^T G(t) dt \ll T^{5/4}.$$

To prove that $b_0 = \pi$ (cf. (2.4)) we note that, using the definition of $E(T)$ and applying integration by parts,

$$\begin{aligned} (2.9) \quad L_1\left(\frac{1}{T}\right) &= \int_0^\infty e^{-t/T} \left(\log\left(\frac{t}{2\pi}\right) + 2\gamma + E'(t) \right) dt \\ &= T \int_0^\infty e^{-x} \left(\log x + \log \frac{T}{2\pi} + 2\gamma \right) dx + \frac{1}{T} \int_0^\infty E(t) e^{-t/T} dt \\ &= T \left(\log\left(\frac{T}{2\pi}\right) + \gamma \right) + \frac{1}{T^2} \int_0^\infty \int_0^t E(u) du \cdot e^{-t/T} dt \\ &= T \left(\log\left(\frac{T}{2\pi}\right) + \gamma \right) + \frac{1}{T^2} \int_0^\infty (\pi t + O(t^{3/4})) e^{-t/T} dt \\ &= T \left(\log\left(\frac{T}{2\pi}\right) + \gamma \right) + \pi + O(T^{-1/4}), \end{aligned}$$

where we also used the O -bound of (2.7) and

$$\int_0^\infty e^{-x} \log x dx = \Gamma'(1) = -\gamma.$$

A comparison of (2.3) and (2.9) proves that $b_0 = \pi$, as asserted by (2.4).

We return now to the proof of Theorem 1. Let

$$(2.10) \quad \bar{L}_k(s) := \int_1^\infty |\zeta(\tfrac{1}{2} + iy)|^{2k} e^{-ys} dy \quad (k \in \mathbb{N}, \sigma = \Re s > 0).$$

Then we have by absolute convergence, taking σ sufficiently large and making the change of variable $xy = t$,

$$\begin{aligned} (2.11) \quad \int_0^\infty \bar{L}_k(x) x^{s-1} dx &= \int_0^\infty \left(\int_1^\infty |\zeta(\tfrac{1}{2} + iy)|^{2k} e^{-yx} dy \right) x^{s-1} dx \\ &= \int_1^\infty |\zeta(\tfrac{1}{2} + iy)|^{2k} \left(\int_0^\infty x^{s-1} e^{-xy} dx \right) dy \\ &= \int_1^\infty |\zeta(\tfrac{1}{2} + iy)|^{2k} y^{-s} dy \int_0^\infty e^{-t} t^{s-1} dt = \mathcal{Z}_k(s) \Gamma(s). \end{aligned}$$

Further we have

$$\begin{aligned} \int_0^\infty \bar{L}_1(x) x^{s-1} dx &= \int_0^1 \bar{L}_1(x) x^{s-1} dx + \int_1^\infty \bar{L}_1(x) x^{s-1} dx \\ &= \int_1^\infty \bar{L}_1(1/x) x^{-1-s} dx + A(s) \quad (\sigma > 1), \end{aligned}$$

where

$$A(s) := \int_1^\infty \bar{L}_1(x) x^{s-1} dx$$

is an entire function. Since

$$\bar{L}_1(1/x) = L_1(1/x) - \int_0^1 |\zeta(\tfrac{1}{2} + iy)|^2 e^{-y/x} dy \quad (x \geq 1),$$

it follows from (2.11) with $k = 1$ by analytic continuation that, for $\sigma > 1$,
(2.12)

$$\begin{aligned} \mathcal{Z}_1(s) \Gamma(s) &= \int_1^\infty L_1(1/x) x^{-1-s} dx - \int_1^\infty \left(\int_0^1 |\zeta(\tfrac{1}{2} + iy)|^2 e^{-y/x} dy \right) x^{-1-s} dx + A(s) \\ &= I_1(s) - I_2(s) + A(s), \end{aligned}$$

say. Clearly for any integer $M \geq 1$

$$\begin{aligned} (2.13) \quad I_2(s) &= \int_1^\infty \int_0^1 |\zeta(\tfrac{1}{2} + iy)|^2 \left(\sum_{m=0}^M \frac{(-1)^m}{m!} \left(\frac{y}{x} \right)^m + O_M(x^{-M-1}) \right) dy x^{-1-s} dx \\ &= \sum_{m=0}^M \frac{(-1)^m}{m!} h_m \cdot \frac{1}{m+s} + H_M(s), \end{aligned}$$

say, where $H_M(s)$ is a regular function of s for $\sigma > -M - 1$, and

$$h_m := \int_0^1 |\zeta(\tfrac{1}{2} + iy)|^2 y^m dy$$

is a constant. Inserting (2.3) in $I_1(s)$ in (2.12) we have, for $\sigma > 1$,

$$\begin{aligned} (2.14) \quad I_1(s) &= \int_1^\infty \left(\log \frac{x}{2\pi} + \gamma \right) \sum_{n=0}^N a_n x^{-2n-s} dx + \int_1^\infty \sum_{n=0}^N b_n x^{-1-2n-s} dx + K_N(s) \\ &= \sum_{n=0}^N a_n \left(\frac{1}{(2n+s-1)^2} + \frac{\gamma - \log 2\pi}{2n+s-1} \right) + K_N(s), \end{aligned}$$

say, where $K_N(s)$ is regular for $\sigma > -2N$. Taking $M = 2N$ it follows from (2.12)–(2.14) that

$$\begin{aligned} (2.15) \quad \mathcal{Z}_1(s) \Gamma(s) &= \sum_{n=0}^N a_n \left(\frac{1}{(2n+s-1)^2} + \frac{\gamma - \log 2\pi}{2n+s-1} \right) \\ &\quad + \sum_{m=0}^{2N} \frac{(-1)^m}{m!} h_m \cdot \frac{1}{m+s} + R_N(s), \end{aligned}$$

say, where $R_N(s)$ is a regular function of s for $\sigma > -2N$. This holds initially for $\sigma > 1$, but by analytic continuation it holds for $\sigma > -2N$. Since N is arbitrary and $\Gamma(s)$ has no zeros, it follows that (2.15) provides meromorphic continuation of $\mathcal{Z}_1(s)$ to $\mathbb{C}$. Taking into account that $\Gamma(s)$ has simple poles at $s = -m$ ($m = 0, 1, 2, \dots$) with residues $\frac{(-1)^m}{m!}$, and that near $s = 1$ we have the Taylor expansion

$$\frac{1}{\Gamma(s)} = 1 + \gamma(s-1) + \sum_{n=2}^{\infty} \frac{f_n}{n!} (s-1)^n$$

with $f_n = (1/\Gamma(s))^{(n)} \Big|_{s=1}$, it follows that

$$\begin{aligned} \mathcal{Z}_1(s) &= \frac{1}{(s-1)^2} + \frac{2\gamma - \log(2\pi)}{s-1} \\ (2.16) \quad &+ \frac{1}{\Gamma(s)} \left\{ \sum_{n=1}^N a_n \left(\frac{1}{(2n+s-1)^2} + \frac{\gamma - \log 2\pi}{2n+s-1} \right) \right\} \\ &+ \frac{1}{\Gamma(s)} \left(\sum_{m=0}^{2N} \frac{(-1)^m}{m!} h_m \cdot \frac{1}{m+s} \right) + U_N(s), \end{aligned}$$

say, where $U_N(s)$ is a regular function of s for $\sigma > -2N$. Moreover, the function $1/(\Gamma(s)(2n+s-1))$ is regular for $n \in \mathbb{N}$ and $s \in \mathbb{C}$. Therefore (2.16) provides analytic continuation to $\mathbb{C}$, showing that besides $s = 1$ the only poles of $\mathcal{Z}_1(s)$ can be simple poles at $s = 1 - 2n$ for $n \in \mathbb{N}$, as asserted by Theorem 1. With more care the residues at these poles could be explicitly evaluated.

3. Proof Theorem 2

We suppose that $X \gg t$ and start from

$$\mathcal{Z}_1(s) = \left(\int_1^X + \int_X^\infty \right) |\zeta(\tfrac{1}{2} + ix)|^2 x^{-s} dx \quad (\sigma = \Re s > 1).$$

We use $|\zeta(\tfrac{1}{2} + ix)|^2 = \log(x/2\pi) + 2\gamma + (E(x) - \pi)'$ and integrate by parts to obtain

$$\begin{aligned} \mathcal{Z}_1(s) &= \int_1^{t^{1-\varepsilon}} |\zeta(\tfrac{1}{2} + ix)|^2 x^{-s} dx + \int_{t^{1-\varepsilon}}^X |\zeta(\tfrac{1}{2} + ix)|^2 x^{-s} dx \\ (3.1) \quad &+ \frac{X^{1-s}}{s-1} \left(\frac{1}{s-1} + \log X + 2\gamma - \log 2\pi \right) \\ &- (E(X) - \pi) X^{-s} + s \int_X^\infty (E(x) - \pi) x^{-s-1} dx \\ &= I_1 + I_2 + O(X^{-\sigma} + t^{-1} X^{1-\sigma} \log X) + I_3, \end{aligned}$$

provided $E(X) = 0$. But as proved in [6], every interval $[T, T + C\sqrt{T}]$ with sufficiently large $C > 0$ contains a zero of $E(t)$, hence we can assume that $E(X) = 0$ is fulfilled with suitable X . Since $E(x)$ is $\asymp x^{1/4}$ in mean square (see e.g., [5] or [7]), it follows by analytic continuation that (3.1) is valid for $\sigma > \frac{1}{4}$, $t \geq t_0 > 0$, $t \leq X \leq t^A$ ($A \geq 1$). By repeated integration by parts it is found that $I_1 \ll t^{-1}$. The integral I_2 is split in $O(\log T)$ subintegrals of the form

$$J_K = \int_{K/2}^{5K/2} \varphi(x) |\zeta(\tfrac{1}{2} + ix)|^2 x^{-s} dx,$$

where $\varphi(x) \in C^\infty$ is a nonnegative, smooth function supported in $[K/2, 5K/2]$ that is equal to unity in $[K, 2K]$. To bound J_K one can start from (1.1) with $k = 1$ and use the Mellin inversion formula (see [10] for a detailed discussion concerning the analogous situation with the fourth moment of $|\zeta(\frac{1}{2} + it)|$), which yields

$$|\zeta(\tfrac{1}{2} + ix)|^2 = \frac{1}{2\pi i} \int_{(1+\varepsilon)} \mathcal{Z}_1(s) x^{s-1} ds \quad (x > 1).$$

Here we replace the line of integration by the contour $\mathcal{L}$, consisting of the same straight line from which the segment $[2 + \varepsilon - i, 1 + \varepsilon + i]$ is removed and replaced by a circular arc of unit radius, lying to the left of the line, which passes over the pole $s = 1$ of the integrand. By the residue theorem we have

$$|\zeta(\tfrac{1}{2} + ix)|^2 = \frac{1}{2\pi i} \int_{\mathcal{L}} \mathcal{Z}_1(s) x^{s-1} ds + \log\left(\frac{x}{2\pi}\right) + 2\gamma \quad (x > 1).$$

This gives

$$J_K = \frac{1}{2\pi i} \int_{\mathcal{L}} \mathcal{Z}_1(w) \int_{K/2}^{5K/2} \varphi(x) x^{w-s-1} dx dw + O(t^{-1}).$$

Repeated integration by parts in the x -integral shows that only the portion $|\Im w - \Im s| \leq t^\varepsilon$ gives a non-negligible contribution. Hence replacing $\mathcal{L}$ by the line $\Re w = \frac{1}{2}$ we obtain

$$(3.2) \quad J_K \ll_\varepsilon K^{\frac{1}{2}-\sigma} t^\varepsilon \max_{t-t^\varepsilon \leq v \leq t+t^\varepsilon} |\mathcal{Z}_1(\tfrac{1}{2} + iv)| + t^{-1}.$$

To estimate I_3 in (3.1) we integrate by parts and use (2.7) and (2.8). We obtain

$$\begin{aligned} \int_X^\infty (E(x) - \pi) x^{-s-1} dx &= -G(X) X^{-s-1} + (s+1) \int_X^\infty G(x) x^{-s-2} dx \\ &= O(X^{-1/4-\sigma}) - (s+1) G_1(X) X^{-s-2} + (s+1)(s+2) \int_X^\infty G_1(x) x^{-s-3} dx \\ &\ll X^{-1/4-\sigma} + t^2 X^{-3/4-\sigma}. \end{aligned}$$

Thus it follows from (3.1) and (3.2) that we have, for $\frac{1}{2} \leq \sigma \leq 1$, $t \geq t_0$,

$$(3.3) \quad \begin{aligned} \mathcal{Z}_1(\sigma + it) &\ll_{\varepsilon} t^{\frac{1}{2}-\sigma+\varepsilon} \max_{t-t^{\varepsilon} \leq v \leq t+t^{\varepsilon}} |\mathcal{Z}_1(\tfrac{1}{2} + iv)| + (X^{1-\sigma}t^{-1} + t^{-1}) \log t \\ &\quad + X^{-\sigma} + tX^{-1/4-\sigma} + t^3X^{-3/4-\sigma}. \end{aligned}$$

In (3.3) we choose X to satisfy

$$X^{1-\sigma}t^{-1} = t^3X^{-3/4-\sigma}, \quad \text{i.e.,} \quad X = t^{16/7}.$$

Then $tX^{-1/4-\sigma} + X^{-\sigma} \ll t^3X^{-3/4-\sigma}$, and (1.3) follows from (3.3).

In conclusion, one may ask what should be the true order of magnitude of $\mathcal{Z}_1(\sigma + it)$. This is a difficult question, but it seems reasonable to expect that

$$(3.4) \quad \mathcal{Z}_1(\sigma + it) \ll_{\varepsilon} t^{\frac{1}{2}-\sigma+2\mu(\frac{1}{2})+\varepsilon} \quad (\tfrac{1}{2} \leq \sigma \leq 1, t \geq t_0 > 0)$$

holds, where

$$\mu(\sigma) := \limsup_{t \rightarrow \infty} \frac{\log |\zeta(\sigma + it)|}{\log t}$$

is the Lindelöf exponent of $\zeta(s)$. One could try to obtain (3.4) from (1.3) by refining the estimation of $\mathcal{Z}_1(s)$ in [11]. This procedure leads to an exponential sum with the divisor function which is estimated as $\ll_{\varepsilon} t^{2\mu(\frac{1}{2})+\varepsilon}$ (this, of course, is non-trivial and needs elaboration).

REFERENCES

- [1] F.V. Atkinson, The mean value of the zeta-function on the critical line, *Quart. J. Math. Oxford* **10**(1939), 122-128.
- [2] F.V. Atkinson, The mean value of the zeta-function on the critical line, *Proc. London Math. Soc.* **47**(1941), 174-200.
- [3] J.B. Conrey, D.W. Farmer, J.P. Keating, M.O. Rubinstein and N.C. Snaith, Integral moments of L -functions, 71pp, [ArXiv.math.NT/0206018](https://arxiv.org/abs/math/0206018).
- [4] J.L. Hafner and A. Ivić, On the mean square of the Riemann zeta-function on the critical line, *J. Number Theory* **32**(1989), 151-191.
- [5] A. Ivić, The Riemann zeta-function, *John Wiley and Sons*, New York, 1985.
- [6] A. Ivić, Large values of certain number-theoretic error terms, *Acta Arith.* **56**(1990), 135-159.
- [7] A. Ivić, Mean values of the Riemann zeta-function, LN's **82**, *Tata Institute of Fundamental Research*, Bombay, 1991 (distr. by Springer Verlag, Berlin etc.).
- [8] A. Ivić, On the fourth moment of the Riemann zeta-function, *Publs. Inst. Math. (Belgrade)* **57(71)** (1995), 101-110.
- [9] A. Ivić, The Laplace transform of the fourth moment of the zeta-function, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.* **11**(2000), 41-48.

- [10] A. Ivić, On some conjectures and results for the Riemann zeta-function, *Acta. Arith.* **99**(2001), 115-145.
- [11] A. Ivić, M. Jutila and Y. Motohashi, The Mellin transform of powers of the Riemann zeta-function, *Acta Arith.* **95**(2000), 305-342.
- [12] M. Jutila, Mean values of Dirichlet series via Laplace transforms, in “*Analytic Number Theory*” (ed. Y. Motohashi), London Math. Soc. LNS **247**, *Cambridge University Press*, Cambridge, 1997, 169-207.
- [13] M. Jutila, The Mellin transform of the square of Riemann's zeta-function, *Periodica Math. Hung.* **42**(2001), 179-190.
- [14] H. Kober, Eine Mittelwertformel der Riemannschen Zetafunktion, *Compositio Math.* **3**(1936), 174-189.
- [15] Y. Motohashi, A relation between the Riemann zeta-function and the hyperbolic Laplacian, *Annali Scuola Norm. Sup. Pisa, Cl. Sci. IV ser.* **22**(1995), 299-313.
- [16] Y. Motohashi, Spectral theory of the Riemann zeta-function, *Cambridge University Press*, 1997.
- [17] E.C. Titchmarsh, The Theory of the Riemann Zeta-Function (2nd ed.), *Clarendon Press*, Oxford, 1986.

ALEKSANDAR IVIĆ
KATEDRA MATEMATIKE RGF-A
UNIVERSITET U BEOGRADU
ĐUŠINA 7, 11000 BEOGRAD
SERBIA AND MONTENEGRO
E-MAIL: aivic@matf.bg.ac.yu,
ivic@rgf.bg.ac.yu